

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : A61K 31/725</p>	<p>A1</p>	<p>(11) International Publication Number: WO 94/23725 (43) International Publication Date: 27 October 1994 (27.10.94)</p>
<p>(21) International Application Number: PCT/CA94/00207 (22) International Filing Date: 15 April 1994 (15.04.94) (30) Priority Data: 2,094,203 16 April 1993 (16.04.93) CA (71) Applicant (for all designated States except US): NOR-PHARMCO INC. [CA/CA]; 890 Yonge Street, Second floor, Toronto, Ontario M4W 3P4 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): WILLOUGHBY, Derek, A. [GB/GB]; The Medical College of St. Bartholomew's Hospital, Charterhouse Square, London EC1M 6BQ (GB). ALAM, Chandan [GB/GB]; The Medical College of St. Bartholomew's Hospital, Charterhouse Square, London EC1M 6BQ (GB). ASCULAI, Samuel, Simon [US/CA]; Apartment #Th13, 53 McCaul Street, Toronto, Ontario M6M 2B6 (CA). FALK, Rudolf, Edgar [CA/CA]; 40 Riverview Drive, Toronto, Ontario M4N 3C7 (CA). HARPER, David, William [CA/CA]; 210 Morrison Road, Oakville, Ontario L6J 4J4 (CA). (74) Agents: HUGHES, Ivor, M. et al.; Suite 200, 175 Commerce Valley Drive West, Thornhill, Ontario L3T 7P6 (CA).</p>		<p>(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KG, KP, KR, KZ, LK, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.</p>
<p>(54) Title: COMPOSITIONS, FOR INHIBITION, CONTROL AND REGRESSION OF ANGIOGENESIS, CONTAINING HYALURONIC ACID AND NSAID (57) Abstract The use of: (a) a non-steroidal anti-inflammatory agent, and (b) hyaluronic acid and/or salts thereof and/or homologues, analogues, derivatives, complexes, esters, fragments, and subunits of hyaluronic acid, in the manufacture of a pharmaceutical composition for inhibiting, controlling and/or regressing angiogenesis in a therapy wherein dosage amounts taken from the composition each comprise: (1) a therapeutically effective amount of component (a); and (2) a therapeutically effective amount of the hyaluronic acid and/or salts thereof and/or homologues, analogues, derivatives, complexes, esters, fragments, and sub-units of hyaluronic acid, the pharmaceutical composition being characterized in that for each dosage amount taken from the pharmaceutical composition, the amount of components (a) and (b) inhibit, control and/or regress angiogenesis.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Compositions, for inhibition, control and regression of angiogenesis, containing hyaluronic acid and NSAID.

FIELD OF INVENTION

5 This invention relates to the inhibition, control and regression of angiogenesis and finds one particular application in the inhibition of angiogenesis in cancer treatment for example as an adjuvant to known cancer treatments for prevention of metastasis.

BACKGROUND OF THE INVENTION

10 In an article entitled "Solid cores of tumors keeping out best drugs" by Sandra Blakeslee published in the July 8, 1989 edition of the Globe and Mail, Toronto, Ontario, Ms. Blakeslee submitted that a growing number of researchers believe that a basic misunderstanding of the
15 structure of solid tumors has led researchers into designing cancer drugs that are doomed to fail in many patients.

She relates that, Dr. Herberman, Director of the Pittsburgh Cancer Center, said that for decades, cancer
20 researchers have simply developed drugs, put them in the bloodstream and assumed they would be carried to the tumor giving almost no consideration to how uniformly the drug is distributed once it reaches the tumor.

Her article also provided that according to Dr.
25 Judah Folkman, a leading researcher on blood growth factors at the Harvard Medical School, for a long time, physicians have been taught that tumors outgrow their blood supply. According to the article that statement is not true. Tumors compress their blood supply. This
30 compression makes it harder to administer drugs.

The article provides further that most people think a tumor is nothing but a collection of cancer cells. According to Dr. Jain, another researcher, in reality the tumor is only 50 per cent cells. The other half is blood
35 vessels and interstitial space. Interstitial space in a tumor, he said, can be likened to the space between marbles packed in a box.

The article further provides that no matter how biological agents are mixed and administered, they must

cross a blood-vessel wall and then cross the interstitium to reach their targets, cancer cells. The article continues that every tumor is different and there are different regions within each. Moreover, tumors change
5 daily as they grow and rearrange parts. Most blood vessels inside tumors are highly disorganized as they take tortuous turns and grow peculiar attachments to nearby vessels.

In general, Dr. Jain said, as a tumor grows, its
10 outer region recruits new blood vessels from surrounding normal tissue. (The tumour needs a blood supply to grow; metastasis also needs a blood supply to develop). The tumour also forms several abnormal blood vessels of its own. As the tumor grows in a confined space, many of the
15 twisted blood vessels near its center are crushed. In turn, the tumor cells near them appear to die, although they grow into active cancer if transplanted into other animals. High pressures build up in these necrotic regions. Both blood vessels and liquid plasma in the
20 interstitial spaces are squeezed. The pressure, therefore, prevents blood-borne molecules, including oxygen, from entering the central tumor areas.

Pressure is not uniform in normal tissue, Dr. Jain said, so a large molecule such as an antibody would
25 reach its target through convection induced by pressure differences. But in the center of a tumor, pressure is uniformly high, blocking convection.

Molecules also migrate by diffusion Dr. Jain said, which is similar to the way a drop of ink spreads in
30 water.

But he indicated that he measured antibody diffusion in tumours and found that it can take days, weeks or months for such large molecules to reach uniform concentration by diffusion in tumours. By then, it may be
35 too late for treatments to do any good.

European Patent Application 0295092 purports to teach a vehicle together with fragments of hyaluronic acid for delivering of the fragments of hyaluronic acid into

the skin to reach the dermal layer of the skin to increase the development of blood vessels for stimulating hair growth or regrowth. The preferred fragments of hyaluronic acid are polysaccharides containing from 7 to 25 monosaccharide units. The patent provides it is apparent that the larger the fragments of hyaluronic acid, the greater the difficulty there is in delivering the fragments to the dermal layer of the skin, unless there is also present in the composition a means for enhancing the activity of said fragments.

The combination may thus include a means for enhancing the activity of the fragments of hyaluronic acid especially to improve their penetration through the skin following topical application. Some activity enhancers, it is alleged, also function as vehicles for the fragments of the hyaluronic acid.

Some activity enhancers are also alleged to possess the ability to stimulate or increase hair growth. Minoxidil is asserted among others to be such an activity enhancer. Thus both the fragments of hyaluronic acid and minoxidil are alleged to stimulate hair growth both delivered by a vehicle.

There have been extensive studies to determine the defect in immune function that allows a tumor cell to develop. It was postulated initially by Jerne, and subsequently by Burnett that the immune system's major role was that of immunological surveillance to destroy abnormal cells. The concept of surveillance, while somewhat simplistic, remains an accepted concept for the elaborate mechanism of immune recognition and function that is present in the higher species - mammals.

It has then been postulated that tumors develop because of local or generalized immune suppression. However, as pointed out by Moller, if general immune suppression occurs, it is only certain types of neoplastic disorders that develop, mainly those of the lymphoreticular system. This observation is correct and represents a major challenge to the immune surveillance

theory unless a specific reason can be shown as to why the individual cancer cell can develop plus individually evade the immune system.

It was demonstrated experimentally in 1974 that
5 defects of macrophage function may exist in neoplastic disease.

The initial experiments found suppressor cells to be part of the immune system; these were either of the T-cell type of the macrophage cell system. There was
10 presence demonstrated in neoplasia, chronic bacterial infection, recovery from massive injury and chronic fungal infection.

There has been repeated demonstration in experimental animals, that the macrophage cell function is
15 altered in neoplastic disease. The macrophages in the animal's systems appeared "blocked" in their function. Generally when removed from the in vivo situation, washed in saline and cultured, they could perform normally. This block has been shown to be related to the excessive
20 production of prostaglandin by neoplastic tissue or by the macrophage itself.

In the basic research efforts in the latter '70s and the early 80's, there existed considerable confusion as to what role immunotherapy should take in cancer.
25 Activation or "hyping" of macrophages was thought to be important. However, in an examination by Romans and Falk of peritoneal macrophages obtained from patients with neoplastic disease, there was definite evidence that these macrophages were already activated yet were co-existing
30 with cancer cells and not causing their destruction.

In the early part of this year it has been shown by several independent investigators that the malfunction of macrophages or the putitive block is due to excessive prostaglandin and that this can be altered in tissue
35 culture by corticosteroids, ASA, and the non-steroidal anti-inflammatory drugs, i.e. indomethacin, and naproxen (Naprosyn™). Again, in animal tumors it was repeatedly demonstrated that these substances could alter the

response to neoplastic cells and that various combinations of these substances employed with immune enhancing agents could produce very credible success in eliminating experimental tumors. Lala and co-workers combined
5 Indomethacin therapy with Interleukin 2 and showed that this could effect a cure with experiment neoplasm.

There were continued problems with the use of any of these agents in the actual human in vivo experience. All of the non-steroidal anti-inflammatory
10 agents (NSAID) produced major toxicity in terms of gastrointestinal, neurological, and other areas. Thus, the basis of the present approach is that under general circumstances the use of these agents in human disease, in sufficient amounts, the drug will penetrate to any
15 pathological tissue to alter therapeutically local prostaglandin production. While intravenous preparations exist of Indomethacin and now of other agents, the data is overwhelming, as is our own experience, that using these drugs alone produces prohibitive side effects in human
20 subjects. Therefore only insufficient amounts can be brought into the body systemically for example to effect more than occasional responses in neoplasm.

New blood vessel formation is essential for the development of tumours and chronic inflammatory
25 granulomatous tissue such as pannus in rheumatoid arthritis. The restriction of angiogenesis in such disease states is a valid therapeutic target for the restriction of tissue growth.

Sub-retinal neovascularisation is a major
30 ophthalmic problem in the western world today. A small percentage of patients are treated with laser photocoagulation when a sub-retinal new vessel grows from the middle choroidal layer of the eye under the retina, destroying macula or sharp vision. Most vessels grow too
35 close to the center however, and a laser burn would therefore destroy any potential central vision.

Alpha interferon injections three times a week have also been proposed for the treatment of sub-retinal

neovascularisation. The side effects however from the injections especially in elderly people are very significant, being detrimental to the procedure. Additionally, the logistics of arranging the injections from freshly made up preparations are very difficult, to say the least.

In arthritis, a pannus develops. In order for the pannus to develop, vascularization must be present as the pannus develops by the vascularization accompanied by deposition of corrective tissue beneath the corneal epithelium. One result is the overgrowth of connective tissue on the articular surface of the diarthrodial joint.

Applicants have now developed methods, and compositions suitable for use, for the inhibition of angiogenesis (the formation and differentiation of blood vessels) which finds among other applications two applications, the treatment of tumours and as an adjuvant to any cancer treatment for the inhibition of metastasis of the cancer. Applicants have also developed new methods and new compositions suitable for use for the regression of angiogenesis (the regression of blood vessel growth).

Applicant's invention can also be used for the treatment of other diseases and conditions whose successful treatment would benefit from inhibiting, controlling and/or regressing blood vessel growth for example, sub-retinal neovascularisation and the reduction of the effects of arthritis. If a person is prone to arthritis, the administration of pharmaceutical compositions according to Applicants' invention, for example systemically (for example by intravenous administration) may even prevent the arthritis or at least reduce the effects of the arthritis.

It is therefore an object of this invention to provide a method for the inhibition, control and regression of angiogenesis, compositions for the inhibition, control and regression of angiogenesis and the use of such compositions for the inhibition, control and regression of angiogenesis.

Further and other objects of the invention will realized by those skilled in the art from the following summary of the invention and detailed description of embodiments thereof.

5 SUMMARY OF THE INVENTION

According to one aspect of the invention, there is provided a process for the inhibition, control and/or regression of angiogenesis, (for example inhibition of blood vessel growth to a malignant tumour, cutting off
10 blood vessel growth or development, in to a malignant tumour) in a mammal (for example a human), the process comprising the steps of administering an effective dosage amount of a pharmaceutical composition for the inhibition, control and/or regression of angiogenesis to a site on/in
15 the mammal in need of inhibition, control and/or regression of angiogenesis usually over a number of weeks or months on a regular basis, each effective dosage amount of the composition comprising in (a homogeneous admixture) an effective non-toxic dosage amount of an NSAID (non-steroidal anti-inflammatory agent) for example diclofenac
20 (for example dissolved in the composition) and an effective non-toxic dosage amount of hyaluronic acid and/or salt thereof (for example sodium hyaluronate) and/or homologues, analogues, derivatives, complexes, esters, fragments and/or sub-units of hyaluronic acid
25 preferably sodium hyaluronate.

According to an embodiment of the invention the composition may be administered systemically (for example intravenously, intra arterially, intraperitoneally,
30 intrapleurally, by direct injection into for example a tumour, and the like) and may be used as an adjuvant to any cancer treatment.

When an NSAID, for example indomethacin (dissolved in n-methyl glucamine), naproxen, (+/-) tromethamine salts of ketorolac, ibuprofen, piroxicam,
35 propionic acid derivatives, acetylsalicylic acid, flunixin, diclofenac, diclofenac sodium or other NSAID is administered with greater than 200 mg of a form hyaluronic

acid (for example 200-1,000 mg of sodium hyaluronate) for a 70 kg person with the effective amount of the NSAID (in one instance diclofenac), no major toxic side effects occur such as gastro-intestinal distress, neurological abnormalities, depression, etc., even at elevated amounts of the NSAID (if necessary). If the amount of hyaluronic acid is decreased below that amount, side effects may begin to occur.

According to another embodiment the composition may also be administered intradermally, applied topically for delivery into the skin, or administered rectally or put on a patch to be secured to the skin of the patient. Whatever the route of administration, the form is a homogeneous composition whether sterile water containing solution for systemic administration or a cream, lotion or gel for topical administration. Whatever the composition it may be packaged in an appropriate container (intravenous bag, vial for injection, tube or jar of cream).

According to another aspect of the invention the use of a combination of a form of hyaluronic acid (for example hyaluronic acid and/or salt thereof and/or a homologue, analogue, derivative, complex, ester, fragment and/or sub-unit of hyaluronic acid) (for example sodium hyaluronate) and a non-steroidal anti-inflammatory agent (NSAID), for example diclofenac sodium is provided for the inhibition, control and/or regression of angiogenesis.

In such use an effective non-toxic amount of a combination of the form of hyaluronic acid (for example sodium hyaluronate) and NSAID (for example diclofenac sodium) may be used for inhibition, control and regression of angiogenesis.

Thus an effective non-toxic dosage amount of a composition comprising an effective non-toxic dosage amount of sodium hyaluronate and a therapeutically effective non-toxic dosage amount of NSAID for example diclofenac sodium may be used to inhibit angiogenesis. The amount of NSAID (for example diclofenac) administered

systemically may be about 15 to about 100 mg of NSAID for example 35 mg/70 kg person (1/2 mg/kg). The NSAID may be administered in larger amounts provided the amounts are non-toxic, for example to 420 mg/70 kg person (6 mg/kg).

5 Where the NSAID is administered systemically (e.g. by injection, etc.) a guide with respect to the amount of the form of hyaluronic acid is that for every about 15 mg of NSAID, about 50 mg of the form of hyaluronic acid may be used (e.g. sodium hyaluronate).

10 For topical application the amount of the form of diclofenac sodium may be in excess of about 5-10 mg/cm² of skin or exposed tissue in the dosage amount to which the dosage amount of the composition is applied. The form of hyaluronic acid, for example sodium hyaluronate, may
15 also be in excess of about 5 - 10 mg/cm² in each dosage amount.

Thus according to another aspect of the invention a pharmaceutical composition (suitable for systemic, and/or topical application [on the skin,
20 rectally, on the mucosa, etc.]) is provided (for example a multigram pharmaceutical composition for use topically) effective for the inhibition, controlling and regression of angiogenesis, the pharmaceutical composition containing a plurality of dosage amounts for inhibiting, controlling
25 and regression of angiogenesis, each said dosage amount comprising a therapeutically effective non-toxic (to the patient) dosage amount of the NSAID (for example diclofenac) and an effective non-toxic dosage amount of the hyaluronic acid and/or salts thereof and/or
30 homologues, analogues, derivatives, complexes, esters, fragments, and/or subunits of hyaluronic acid effective to inhibit, control and regress angiogenesis.

The pharmaceutical composition may comprise suitable excipients depending upon the route of
35 administration for example sterile water (systemic administration), excipients to make a gel, lotion or cream (topical administration) or whatever may be the route (for example a solubilizer (such as methoxypolyethelene glycol

350) and a preservative such as benzyl alcohol).

According to yet another aspect of the invention, a dosage amount of a pharmaceutical composition is provided for inhibition, controlling and regressing of angiogenesis, the composition comprising:

(1) a non-steroidal anti-inflammatory agent (NSAID) for example diclofenac sodium; and

(2) hyaluronic acid and/or salts thereof (for example sodium hyaluronate) and/or homologues, analogues, derivatives, complexes esters, fragments and subunits of hyaluronic acid

characterized in that said composition:

(a) is in a dosage form (e.g. in a cream, lotion, gel, intravenous solution, injectable, etc.) which is suitable for administration (systemically or topically, etc.); and

(b) is in such an amount and in such form that component (1) is in an effective dosage amount together with component(2) to inhibit, control and regress angiogenesis (for example which inhibits metastasis). The pharmaceutical composition may comprise a plurality of dosage amounts.

According to still another aspect of the invention the use of

(1) a non-steroidal anti-inflammatory agent (NSAID) for example diclofenac sodium; and

(2) hyaluronic acid and/or salts thereof and/or homologues, analogues, derivatives, complexes, esters, fragments and subunits of hyaluronic acid,

in the manufacture of a pharmaceutical composition for use to inhibit, control and regress angiogenesis in mammals (for example in humans) is provided wherein dosage amounts may be taken from the composition and each dosage amount taken comprises:

a therapeutically effective non-toxic dosage amount of each of components (1) and (2) to inhibit, control and regress angiogenesis.

The composition containing the form of

hyaluronic acid and NSAID provides greater inhibition, control and regression (significantly greater inhibition, control and regression) of angiogenesis than a composition comprising a form of hyaluronic acid (for example sodium hyaluronate) only (without an NSAID). Thus, according to another aspect of the invention Applicants have provided similar methods of treatment, pharmaceutical compositions, dosage amounts and uses comprising forms of hyaluronic acid (for example sodium hyaluronate having a molecular weight less than about 750,000 daltons) without the NSAIDS as it did with the forms of hyaluronic acid with the NSAIDS.

The amount of the form of hyaluronic acid (for example sodium hyaluronate) per dosage amount may vary from about 5 mg for human administration as for example topically (5 mg/cm² of skin or exposed tissue) depending on the route of administration to about 1,000 mg/70 kg person depending upon the route of administration. As there is not toxicity, the form of hyaluronic acid can obviously be administered in a dose excess (for example 3000 mg/70 kg person if administered systemically) without any adverse effects. Amounts of the form of hyaluronic acid (for example sodium hyaluronate) to be administered systemically may be about 50 for each 15 mg of NSAID. Preferably the form of hyaluronic acid (for example sodium hyaluronate) administered, has a molecular weight less than about 750,00 daltons (for example about 150,00 to about 225,000 daltons). While higher molecular weights of the hyaluronic acid and forms thereof may be used to inhibit angiogenesis, where the molecular weight of the hyaluronic acid chosen for use is very large, it is preferred that the form of hyaluronic acid is autoclaved, to break down the hyaluronic acid to fragments of lesser molecular weight or if feasible diluted to permit administration and ensure no coagulation (whatever the route of administration). Where the molecular weight of the form of the form of hyaluronic acid being employed is larger, the concentration of the form of the hyaluronic

acid in the composition may be adjusted, for example be reduced (for example to less than about 1 %) dependent on the molecular weight.

One form of hyaluronic acid and/or salts thereof
5 (for example sodium salt) and homologues, analogues, derivatives, complexes, esters, fragments, and sub-units of hyaluronic acid, preferably hyaluronic acid and salts and thereof, suitable for use with Applicant's invention is a fraction supplied by Hyal Pharmaceutical Corporation.
10 One such fraction is a 15 ml vial of Sodium hyaluronate 20 mg/ml (300 mg/vial - Lot 2F3). The sodium hyaluronate is a 2% solution with a mean average molecular weight of about 225,000. The vial also contains water q.s. which is triple distilled and sterile in accordance with the U.S.P.
15 for injection formulations. The vials of hyaluronic acid and/or salts thereof may be carried in a Type 1 borosilicate glass vial closed by a butyl stopper which does not react with the contents of the vial.

The fraction of hyaluronic acid and/or salts
20 thereof (for example sodium salt) and homologues, analogues, derivatives, complexes, esters, fragments, and sub-units of hyaluronic acid, preferably hyaluronic acid and salts thereof, may comprise hyaluronic acid and/or salts thereof having the following characteristics:

25 a purified, substantially pyrogen-free fraction of hyaluronic acid obtained from a natural source having at least one characteristic selected from the group (and preferably all characteristics) consisting of the following:

30 i) a molecular weight within the range of 150,000-225,000;

ii) less than about 1.25% sulphated mucopoly-saccharides on a total weight basis;

35 iii) less than about 0.6% protein on a total weight basis;

iv) less than about 150 ppm iron on a total weight basis;

v) less than about 15 ppm lead on a

total weight basis;

- vi) less than 0.0025% glucosamine;
- vii) less than 0.025% glucuronic acid;
- viii) less than 0.025% N-acetylglucosamine;
- 5 ix) less than 0.0025% amino acids;
- x) a UV extinction coefficient at 257 nm of less than about 0.275;
- xi) a UV extinction coefficient at 280 nm of less than about 0.25; and

- 10 xii) a pH within the range of 7.3-7.9.
- Preferably, the hyaluronic acid is mixed with water and the fraction of hyaluronic acid has a mean average molecular weight within the range of 150,000-225,000. More preferably, the fraction of hyaluronic acid comprises
- 15 at least one characteristic selected from the group (and preferably all characteristics) consisting of the following characteristics:

- i) less than about 1% sulphated mucopolysaccharides on a total weight basis;
 - 20 ii) less than about 0.4% protein on a total weight basis;
 - iii) less than about 100 ppm iron on a total weight basis;
 - iv) less than about 10 ppm lead on a total
 - 25 weight basis;
 - v) less than 0.00166% glucosamine;
 - vi) less than 0.0166% glucuronic acid;
 - vii) less than 0.0166% N-acetylglucosamine;
 - viii) less than 0.00166% amino acids;
 - 30 x) a UV extinction coefficient at 257 nm of less than about 0.23;
 - xi) a UV extinction coefficient at 280 nm of less than 0.19; and
 - xii) a pH within the range of 7.5-7.7
- 35 Applicants also propose to use sodium hyaluronate produced and supplied by LifeCore™ Biomedical, Inc., having the following specifications:

	<u>Characteristics</u>	<u>Specification</u>
	Appearance	White to cream colored particles
	Odor	No perceptible odor
5	Viscosity Average	< 750,000 Daltons
	Molecular Weight	
	UV/Vis Scan, 190-820 nm	Matches reference scan
	OD, 260 nm	< 0.25 OD units
10	Hyaluronidase Sensitivity	Positive response
	IR Scan	Matches reference
	pH, 10 mg/g solution	6.2 - 7.8
	Water	8% maximum
	Protein	< 0.3 mcg/mg NaHy
15	Acetate	< 10.0 mcg/mg NaHy
	Heavy Metals, maximum ppm	
	As Cd Cr Co Cu Fe Pb Hg Ni	
	2.0 5.0 5.0 10.0 10.0 25.0 10.0 10.0 5.0	
	Microbial Bioburden	None observed
20	Endotoxin	< 0.07EU/mg NaHy
	Biological Safety Testing	Passes Rabbit Ocular Toxicity Test

Another form of sodium hyaluronate is sold under the name Hyaluronan HA-M5070 by Skymart Enterprises, Inc. having the following specifications:

Specifications' Test

Results

	Lot No.	HG1004
30	pH	6.12
	Chondroitin Sulfate	not detected
	Protein	0.05%
	Heavy Metals	Not more than 20 ppm
	Arsenic	Not more than 2 ppm
35	Loss on Drying	2.07%
	Residue on Ignition	16.69%
	Intrinsic Viscosity	12.75 dl/s (XW: 679,000)
	Nitrogen	3.14%

- 15 -

Assay	104.1%
Microbiological Counts	80/g
E. coli	Negative
Mold and Yeast	Not more than 50/g

5

Other forms of hyaluronic acid and/or its salts, and homologues, derivatives, complexes, esters, fragments and sub units of hyaluronic acid may be chosen from other suppliers, for example those described in prior art documents provided the form of hyaluronic acid chosen is suitable for transport of the medicine.

The following references teach hyaluronic acid, sources thereof, and processes for the manufacture and recovery thereof which may be suitable.

15 United States Patent 4,141,973 teaches hyaluronic acid fractions (including sodium salts) having:

20 "(a) an average molecular weight greater than about 750,000, preferably greater than about 1,200,000 - that is, a limiting viscosity number greater than about 1400 cm³/g., and preferably greater than about 2000 cm³/g.;

(b) a protein content of less than 0.5% by weight;

25 (c) ultraviolet light absorbance of a 1% solution of sodium hyaluronate of less than 3.0 at 257 nanometers wavelength and less than 2.0 at 280 nanometers wavelength;

30 (d) a kinematic viscosity of a 1% solution of sodium hyaluronate in physiological buffer greater than about 1000 centistokes, preferably greater than 10,000 centistokes;

35 (e) a molar optical rotation of a 0.1 - 0.2% sodium hyaluronate solution in physiological buffer of less than -11 X 10³ degree - cm²/mole (of disaccharide) measured at 220 nanometers;

(f) no significant cellular infiltration of the vitreous and anterior chamber, no flare

in the aqueous humour, no haze or flare in the vitreous, and no pathological changes to the cornea, lens, iris, retina, and choroid of the owl monkey eye when one milliliter of a 1% solution of sodium hyaluronate dissolved in physiological buffer is implanted in the vitreous replacing approximately one-half the existing liquid vitreous, said HUA being

5 (g) sterile and pyrogen free and

10 (h) non-antigenic."

Canadian Letters Patent 1,205,031 (which refers to United States Patent 4,141,973 as prior art) refers to hyaluronic acid fractions having average molecular weights of from 50,000 to 100,000; 250,000 to 350,000; and 500,000 to 730,000 and discusses processes of their manufacture.

15

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1 comprises two Figures 1(a) and 1(b) which are bar graphs illustrating the effect of formulations on the vasculature within the murine chronic granulomatous air pouch.

20

FIGURE 2 illustrates graphically the results of tests relating to Anti-Angiogenesis: VI Ratio.

FIGURE 3 illustrates graphically the results of tests relating to Anti-Angiogenesis: Carmine.

FIGURE 4 illustrates graphically the results of tests relating to Anti-Angiogenesis: Dry Weight Granuloma.

25

DETAILED DESCRIPTION OF EMBODIMENTS

The invention will now be illustrated with reference to the following experimental data and tests performed.

30

The Applicants chose a simple model of murine angiogenesis induced with a chronic granulomatous reaction to Freund's complete adjuvant in croton oil. The Applicants assessed this process using a modification of this method by the formation of an intravascular cast incorporating carmine red. Chronic granulomatous air pouches were induced by the sc injection of 3 ml air into anaesthetized mice (25-30 g, Tuck Original) and 0.5 ml

35

Freund's complete adjuvant with 0.1% croton oil 24 hours later. The mice were dosed for 6 days and the vascular content assessed by the formation of a vascular cast. This was formed by the intravenous injection of 1 ml 25% carmine red in 10% gelatin at 40°C into warmed mice. This overcame any pharmacological or temperature-related alterations in peripheral vasomotor tone which may have invalidated the results. The carcasses were chilled and the granulomatous air pouch linings dissected. These were dried at 56°C, weighed, and papain digested. The dye was then be dissolved by the addition of 1 ml 0.05 M NaOH, and the samples were then centrifuged at 2500 g for 20 minutes. After these were filtered, the absorbances were read at 490 nm using a multiwell plate reader (Biotek). The results were then expressed as either µg dye/mg dry weight of tissue.

The Applicants have tested whether the topical application of diclofenac/hyaluronan (HA) - e.g. sodium hyaluronate whose molecular weight was less than 750,000 daltons, would have angiostatic activity and whether the HA e.g. sodium hyaluronate alone would have angiostatic static activity.

Topical applications of 0.1 ml hyaluronan and diclofenac (6 mg/kg) were made daily to the surface of the depilated air pouch. Diclofenac, when administered alone, was given with carboxymethylcellulose. The intra-lesional application was carried out by injection into the air pouch, diclofenac alone being administered in sterile saline. On the seventh day after induction the vascularity of the tissue was assessed. Figures 1(a) and 1(b) show that both the topical and intra-lesional administration of hyaluronan (HA) alone and diclofenac (diclo) alone exhibited no significant effect on the tissue vascularity. However, the combination of hyaluronan (HA) and diclofenac (diclo) produced a significant reduction in vascularity. (Figures 1(a) and 1(b) illustrate the effect of hyaluronan and diclofenac, alone or in combination on the development of the

vasculature within the murine chronic granulomatous air pouch. Topical applications of 0.1 ml hyaluronan and diclofenac (6 mg/kg) were made daily to the surface of the depilated air pouch ether alone or in combination (Figure 1(a)). Diclofenac alone was given in carboxymethylcellulose as vehicle. Intra-lesional application was carried out by injection into the air pouch (Figure 1(b)). In this instance diclofenac was given in saline. On the seventh day the vascularity of the tissue was assessed)

**PROTOCOL FOR TOPICAL AND INTRA-POUCH
ADMINISTRATION OF HYALURONIC ACID AND DICLOFENAC
IN COMBINATION**

Introduction:

The formation of a subcutaneous air pouch in the dorsum of mice allows the formation of a lining which responds to produce a chronic inflammatory lesion in response to various antigens, irritants and foreign bodies. It can also be used for the introduction of drugs and various other treatments into the site of inflammation and the collection of inflammatory exudate.

A unique and simple technique has been perfected which quantitatively assesses angiogenesis in the developing inflammatory air pouch, by making a vascular cast incorporating carmine which can then be spectrophotometrically assayed.

Method:

For this experiment female mice (T0, 25-30g, 10 per group) were lightly anesthetized with hypnorm/hypnoval. Air pouches were formed by the subcutaneous injection of 3 ml of filter-sterilized air, into the dorsum of each mouse. After 24 hours, chronic inflammation was induced in the air pouch lining by the injection of 0.5 ml Freund's complete adjuvant (FCA) supplemented with 0.1% croton oil.

Dosing Schedule:

Animals were dosed daily from the time of injection of FCA/croton oil, for 6 Days. At which point the analysis was performed.

5

1. Topical: (See Table One: Topical Application)

Four groups, - 0.1 ml Aqueous Cream * (Thornton & Ross Ltd.)

- 10 - 0.1 ml Aqueous Cream + 6 mg/kg Diclofenac
(HPC lot.9113003)
- 0.1 ml Hyaluronic Acid (sodium
hyaluronate)
(Hyal Pharmaceutical Corporation (HPC))
lot.OG019
- 15 - 0.1 ml Hyaluronic Acid (1% solution sodium
hyaluronate, M.W. less than 750,000
daltons -
e.g. 225,000 daltons)
- 20 + 6 mg/kg Diclofenac

Before the topical application the hair on the dorsum was removed using hair clippers and depilatory cream (Louis Marcel). The skin surface was broken in one or two mice in each group, while using the electric
25 clippers. These animals were discarded from the results.

2. Intra-Pouch: (Injection)

- Four groups, - 0.1 ml Sterile Saline (0.9%)
- 0.1 ml Sterile Saline + 6 mg/kg Diclofenac
(HPC lot.9113003)
- 30 - 0.1 ml Hyaluronic Acid (sodium
hyaluronate)
(HPC lot.OG019)
- 0.1 ml Hyaluronic Acid (1% solution, M.W.
less than 750,000 daltons - e.g. 225,000
daltons) + 6 mg/kg Diclofenac
- 35

Vascular Casting:

Mice were anesthetized with hypnorm/hypnoval and placed on a heated operating platform maintained at 37°C

for 20 minutes. Each mouse was then placed into a water jacketed incubation chamber at 37°C and injected i.v. with 1 ml of 15% carmine dye in 10% gelatin (in Hanks balanced salt solution) with syringe and solution prewarmed to 5 40°C. Cadavers were then chilled at 4°C for 4 hours.

Analysis

The granulomatous tissue was dissected free and dried in an oven at 56°C for 48 hours. The dried granuloma was weighed and then digested in 9 ml of papain 10 solution (12-Units/ml in 0.05 M phosphate buffer, pH 7.0, supplemented with 0.33g/l N-acetyl cysteine), at 56°C for 48 hours. The digests were then made up to 10 ml with 5.0M NaOH and vortexed, to solubilise the dye and centrifuged. The digests were filtered through 0.2µm 15 cellulose nitrate membranes and 200 µl samples aliquoted into 96 well plates and analyzed for dye content by spectrophotometric analysis at 490 nm using a microplate reader. Vascular volume was calculated as carmine content per mg dry mass of tissue.

20 *Commercially available water based cream.

TABLE ONE: Topical Application

					Hyaluronic acid (0.1ml) (1% solution (10 mg/ml), M.W. less than 750,000 daltons)			
Aqueous cream (0.1ml)								
Animal	Dry wt (mg)	Absorbness	mg carmine	ug/dye/mg	Dry wt (mg)	Absorbous	ug carmine	ug dye mg
1	174	.0183	65.5	.376	187	0657	246	1.316
2	303	0.848	319	1.053	178	0.528	197	1.107
3	133	0.613	229	1.722	146	0.148	52.2	0.358
4	163	0.769	289	1.773	190	0.433	161	0.847
5								
6	221	1.014	382	1.728	181	0.505	188	1.039
7	130	0.689	258	1.985	179	0.494	184	1.028
8	122	0.606	227	1.661	160	0.567	212	1.325
9					192	0.631	263	1.229
10	191	0.413	156	0.817				
MEAN	179.6		240.7	1.389	176.62		184.5	1.031
S.e.m.	21.26		346	0.201	561		31.3	0.111
P=					NS		NS	NS

Hyaluronic acid (1% solution) + diclo(6mg/kg)					Aqueous cream + diclo(6mg/kg)			
Animal	Dry wt (mg)	Absorbness	mg carmine	ug/dye/mg	Dry wt (mg)	Absorbous	ug carmine	ug dye mg
1	152	0.185	66.2	0.436	126	0.528	218	1.730
2	176	0.125	43.4	0.242	176	0.420	156	0.886
3	201	0.441	164	0.816	140	0.401	148	1.057
4	149	0.461	171	1.148				
5	196	0.578	216	1.102	124	0.510	190	1.532
6					167	0.464	173	1.035
7	88	0.176	62.8	0.714				
8	131	0.195	70.1	0.535	197	0.808	304	1.534
9	135	0.109	37.3	0.276	207	0.854	231	1.551
10					193	0.217	78.2	0.405
Mean	154.6		103.6	0.658	166.3	187.3	1.216	
S.e.m	13.79		24.3	0.123	11.58		23.6	0.158
P=	NS		0.0104*	0.0209*	NS		NS	NS

n = X

(diclo = Diclofenac)

TABLE TWO: Intra - pouch

	Hyaluronic Acid (1% solution) + diclo(6mg/kg)					Hyaluronic Acid (1% solution)			
	Animal	Dry wt (mg)	Absorbness	mg carmine	ug/dye/mg	Dry wt (mg)	Absorbous	ug carmine	ug dye mg
5	1	104	0.323	68	0.654	126	0.607	130	1.032
	2.	106	0.293	61.5	0.580	128	0.209	43.3	0.337
	3.	140	0.215	44.6	0.319	142	0.710	152	1.069
	4.	133	0.305	64.1	0.482	108	0.603	129	1.194
	5	115	0.460	97.7	0.846	118	0.414	87.8	0.747
10	6	90	0.258	53.9	0.602	97	0.293	61.8	0.637
	7	262	0.271	56.8	0.217	134	0.190	39.2	0.294
	8	72	0.178	36.6	0.509	160	0.494	105	0.654
	9	64	0.171	35.1	0.545	147	0.495	105	0.716
	10	85	0.210	43.5	0.512	130	0.318	66.9	0.517
15	MEAN	117.1		56.2	0.5266	129		91.97	0.7197
	S.e.m.	17.8		5.85	0.0546	5.85		12.24	0.0957
	p=	0.0454*		0.0029**	0.0260*	NS		NS	NS
	Saline					Saline + diclo(6mg/kg)			
	Animal	Dry wt (mg)	Absorbness	mg carmine	ug/dye/mg	Dry wt (mg)	Absorbous	ug carmine	ug dye mg
20	1	211	0.919	197	0.932	139	0.773	166	1.194
	2	98	0.590	126	1.282	158	0.242	50.5	0.320
	3	135	0.390	82.6	0.614	99	0.730	156	1.577
	4	161	0.880	101	1.184	99	0.400	104	1.050
	5	117	0.482	103	0.880	129	0.409	86.7	0.674
25	6	134	0.587	125	0.932	124	0.262	54.8	0.442
	7	138	0.447	94.9	0.686	162	0.402	85.2	0.526
	8	233	0.398	84.3	0.361	135	0.303	63.7	0.471
	9	109	0.438	93	0.854	170	0.468	99.5	0.586
	10	168	0.220	45.7	0.272	122	0.292	61.3	0.520
30	Mean	150.4		114.3	0.7988	133.7		92.75	0.7363
	S.e.m	13.8		15.1	0.1022	7.75		12.77	0.1275
	p=					NS		NS	NS
35									

N = 10

TABLE THREE: Results

	Topical	Carbime dye (ug)			Granuloma dry wt. (mg)			ug dye/mg granuloma		
		Mean	S.e.m.	p=	Mean	S.e.m.	p=	Mean	S.e.m.	p=
5	Aq. cream	240.7	34.6		179.6	21.3		1.389	0.201	
	aq+diclo	187.3	23.6	NS	166.3	11.6	NS	1.216	0.158	NS
	HA	184.3	21.3	NS	176.6	5.6	NS	1.031	0.111	NS
	HA+diclo	103.6	24.3	0.0104*	154.6	13.8	NS	0.658	0.123	0.0209*
	i-pouch									
10	Saline	114.3	15.1		150.4	13.8		0.7988	0.1022	
	Control+	92.6	12.8	NS	133.7	7.8	NS	0.7363	0.1229	NS
	diclo									
	HA	91.9	12.5	NS	129	5.9	NS	0.07197	0.0957	NS
	HA+diclo	56.2	5.85	0.0029**	117.1	17.8	0.0454*	0.5266	0.0546	0.0260*

15 n = 8 for Topical application
 n = 10 for Intra -pouch treatment

20 This data supports the conclusion of HA and Diclofenac, in combination, acting synergistically as an angiostatic agent. Also, it is important to note that although not significant, the trend is such that HA + diclo is more angiostatic than HA alone, the latter being more potent than diclo.

25 All the data has been included for statistical analysis, if however one excludes "obvious" flyers, e.g. table two, animal 7, HA + diclo, the results are more significant.

30 The results of the tests and experiments firmly establish that forms of hyaluronic acid (for example sodium hyaluronate having a molecular weight less than 750,000 daltons - e.g. 225,000 daltons) and NSAID (for example diclofenac) act to inhibit angiogenesis, in for example the granuloma resulting in a reduction in granuloma dry weight.

35 The inhibition of angiogenesis may be used in the treatment and destruction of cancerous tumours. The compositions, dosage amounts taken from the compositions, processes and treatments by the invention may be used as

an adjuvant to any anti-cancer treatment (for example radiation, chemotherapy using anti-cancer drugs, etc.) The invention may also be used to prevent metastasis in cancer patients so that while one tumour is being
5 eradicated, no other malignant tumours develop. Thus the development of tumours is inhibited by inhibition of blood vessel growth to a tumour (cutting off the supply of blood vessels to the tumour). In this regard use of the invention counters, opposes, interferes with and inhibits
10 resulting activity by Tumour Angiogenesis Factor (TAF) produced by a cancerous tumour to increase blood vessel growth to such tumour, thereby inhibiting such growth. A composition, for example comprising sodium hyaluronate and diclofenac administered systemically to a human patient
15 inhibits angiogenesis and the tumour is eradicated. The composition may be administered over a short term or a longer term as required (for example a number of weeks or months as required).

It appears from these initial investigations
20 that the combination of hyaluronan and diclofenac, given either topically or directly into the lesion, results in reduced vascular development during granulomatous inflammation.

FURTHER PROTOCOL

25 In another protocol, the use of the air pouch on the mice was once again employed. Only this time, after formation of the pouches and injection of .5 ml Freund's complete adjuvant (FCA) supplemented with 0.1% croton oil, the granuloma in the mice were permitted to grow for 7
30 days (they were left unhealed for 1 week). The result was the growth of new blood vessels.

Some of the mice were sacrificed at the end of the 7 day period (in the same manner as previously described) while others were treated topically (dosed
35 daily) with:

- 0.1ml aqueous cream (Thornton & Ross Ltd.)
- 0.1ml aqueous cream + 6 mg/kg diclofenac (HPC Lot 9113003)

- 25 -

- 0.1ml hyaluronic acid (HPC Lot 0G019)
- 0.1ml hyaluronic acid (1% solution sodium hyaluronate, molecular weight less than 750,000 daltons - e.g. 225,000 daltons) + 6 mg/kg

5 diclofenac

Some of the mice were sacrificed (in the same manner) after 2 weeks (one week after the commencement of the topical application) and the remainder sacrificed (in the same manner) after 3 weeks (two weeks after the commencement of the topical application).

The carmine red dye displaced the blood in the granulomatous tissue. The granulomatous tissue was dissected and treated as previously described. The weight of the granuloma (including the carmine red dye) and weight of the carmine red dye (which displaced the blood) were determined for each mouse sacrificed at each stage (after 7, 14 and 21 days). The vascularity index (VI) was calculated and the mean vascularity index was calculated for each group as follows:

20

$$\text{Vascularity Index (VI)} = \frac{\text{weight of carmine red dye in GRANULOMA}}{\text{Weight of GRANULOMA (including carmine red dye)}}$$

25

From the following table it can be seen that with both the hyaluronic acid (HA) and HA/diclo combination, the mean dry weight of the granuloma decreased over time. With the HA/diclo composition, the vascularity index (VI) decreased (reduced) over time.. This did not occur with any other composition tested. It is therefore clear, this decrease (reduction) is a measure of the REDUCTION in vascularity (blood supply).

30

TABLE FOUR

The results for all mice sacrificed at one time were dealt with together

		Carminc (ug)	Dry wt. gran. (mg)	VI (ug/mg)
1 week air pouch	mean	173.55	116.71	1.505
	s.e.m.	6.86	4.56	0.062
	p =	--	--	--
2 week aqueous cream	mean	127.87	102.65	1.228
	s.e.m.	17.98	7.33	0.066
	p =	0.0451*	NS	NS
3 week aqueous cream	mean	212.72	116.57	1.822
	s.e.m.	10.66	3.49	0.062
	p =	0.0141*	NS	0.0046**
2 week HA	mean	115.70	101.23	1.136
	s.e.m.	17.12	6.56	0.141
	p =	0.0153	0.0328*	0.0436*
3 week HA	mean	118.68	80.1	1.402
	s.e.m.	14.97	5.44	0.138
	p =	0.0075**	0.0001***	NS
2 week aq. cream/Diclo	mean	108.63	84.41	1.300
	s.e.m.	8.98	5.58	0.089
	p =	0.0084***	0.0006***	NS
3 week aq. cream/Diclo	mean	150.07	112.19	1.353
	s.e.m.	12.07	7.79	0.115
	p =	NS	NS	NS
2 week HA/Diclo	mean	69.63	60.23	1.196
	s.e.m.	8.61	7.72	0.141
	p =	0.0002***	0.0003***	0.0376*
3 week HA/Diclo	mean	66.63	55.31	1.125
	s.e.m.	15.00	8.47	0.212
	p =	0.0002***	0.0002***	0.0224*

s.e.m. = standard error mean

5 p = confidence

For better illustration, the results are shown in the graphs in Figures 2, 3 and 4 which teach

Figure 2 "Anti-angiogenesis": VI Ratio

Figure 3 "Anti-angiogenesis": Carmine

Figure 4 "Anti-angiogenesis": Granuloma

Table Five is also provided setting out the means data in close proximity to one another for ease of comparison and upon which the graphs in Figures 2, 3 and 4 were prepared.

TABLE FIVE

Carmine (ug)			
	Control week 1	week 2	week 3
Aq.Cr. alone	173.55	127.87	212.72
HA alone	173.55	115.7	118.68
Aq. Cr./Diclo	173.55	108.63	150.07
HA/Diclo	173.55	69.63	66.63
dry weight			
granuloma (mg)			
	Control week 1	week 2	week 3
Aq. Cr.	116.71	102.65	116.57
HA alone	116.71	101.23	80.1
Diclo Alone	116.71	84.41	112.19
HA/Diclo	116.71	60.23	55.31
VI (ug/mg)			
	Control week 1	week 2	week 3
Aq. cr.	1.505	1.228	1.822
HA alone	1.505	1.136	1.402
Diclo Alone	1.505	1.3	1.353
HA/Diclo	1.505	1.196	1.125

10 The data with respect to the sodium hyaluronate composition administered to the mice was extrapolated for application to humans. The calculations below are not meant to mean that substantial regression would not be achieved by lesser amounts. In this regard the .1 ml
15 solution of 1% HA solution administered with 6 mg/kg of Diclofenac to the 25-30 gm mice translates to 33.3 mg HA/kg and 6 mg Diclofenac sodium/kg. Thus a 70 kg person would have been administered in excess of 2200 mg of HA and 420 mg of Diclofenac.

It will also be appreciated by those skilled in the art that the processes, uses, compositions and dosage forms according to aspects of the invention may be applied to inhibit angiogenesis in other instances where
5 inhibition of angiogenesis is desired, for example sub-retinal neovascularisation and for the treatment of arthritis or the prevention of further damage thereby including the prevention of the further development of pannus. It is therefore clear that many uses can be made
10 of embodiments and aspects of this invention without departing from the scope thereof. It is therefore intended that all material contained herein be interpreted as illustrative of the invention and not in a limiting sense.

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE AS FOLLOWS:

1. The use of:

(a) a non-steroidal anti-inflammatory agent,

and (b) hyaluronic acid and/or salts thereof and/or homologues, analogues, derivatives, complexes, esters, fragments, and subunits of hyaluronic acid,

in the manufacture of a pharmaceutical composition for inhibiting, controlling and/or regressing angiogenesis in a therapy wherein dosage amounts taken from the composition each comprise:

(1) a therapeutically effective amount of component (1); and

(2) a therapeutically effective amount of the hyaluronic acid and/or salts thereof and/or homologues, analogues, derivatives, complexes, esters, fragments, and sub-units of hyaluronic acid, the pharmaceutical composition being characterized in that for each dosage amount taken from the pharmaceutical composition, the amount of components (1) and (2) inhibit, control and/or regress angiogenesis.

2. The use of Claim 1 wherein component (2) is sodium hyaluronate.

3. The use of Claim 1 wherein the pharmaceutical composition is for sub-retinal neovascularisation.

4. The use of Claim 1 wherein the pharmaceutical composition is for the treatment of arthritis or pannus.

5. The use of Claim 1 wherein the pharmaceutical composition is for the treatment of tumours.

6. The use of Claim 1 wherein the pharmaceutical composition is for an adjunct to cancer treatment.

7. The use of any preceding claim wherein component (2) is sodium hyaluronate having a molecular weight less than about 750,000 daltons.
8. The use of any preceding claim wherein the pharmaceutical composition is for topical application to the skin and/or exposed tissue.
9. The use of Claim 7 wherein the pharmaceutical composition is for systemic administration.
10. The use of Claim 9 wherein the amount of component (2) in each dosage amount exceeds about 50 mg for every 15 mg of NSAID.
11. The use of Claim 9 wherein the form of hyaluronic acid exceeds about 200 mg/70 kg person per dosage amount and the NSAID is between about 15 mg to about 100 mg of NSAID.
12. The use of any preceding claim wherein component (1) the non-steroidal anti-inflammatory drug (NSAID), is selected from diclofenac, diclofenac sodium, indomethacin, naproxen, (+/-) tromethamine salt of ketorolac, ibuprofen, piroxicam, propionic acid derivatives, acetylsalicylic acid and flunixin.
13. The use of:
 - (1) a non-steroidal anti-inflammatory agent (NSAID), and
 - (2) hyaluronic acid and/or salts thereof,in the manufacture of a dosage amount of a pharmaceutical composition for inhibiting, controlling or regressing angiogenesis in humans in a therapy wherein a dosage amount comprises a therapeutically effective amount of said agent (1) and a therapeutically effective amount of the hyaluronic acid and/or salts thereof having a molecular weight less than about 750,000 daltons, the use being characterized in that the amount of components

(1) and (2) is immediately available upon administration to inhibit, control or regress angiogenesis.

14. The use of Claim 10 wherein the dosage amount is for topical application to the skin and/or exposed tissue.

15. The use of Claim 13 wherein the dosage amount is for systemic administration.

16. The use of Claim 12 wherein the amount of component (2) in each dosage amount exceeds about 50 mg for every 15 mg of NSAID.

17. The use of Claim 13 wherein the amount of the form of hyaluronic acid exceeds about 200 mg/70 kg person and the NSAID is between about 15 mg to about 100 mg of NSAID.

18. The use of any preceding claim further comprising pharmaceutically compatible excipients to provide a form for ease of administration depending upon the route of administration.

19. The use of Claim 13 or 14 for administering the composition topically, rectally or to the mucosa.

20. The use of Claim 13 wherein the pharmaceutical composition is for sub-retinal neovascularisation.

21. The use of Claim 13 wherein the pharmaceutical composition is for the treatment of arthritis or pannus.

22. The use of Claim 13 wherein the pharmaceutical composition is for the treatment of tumours.

23. The use of Claim 13 wherein the pharmaceutical composition is for an adjunct to cancer treatment.

24. A method of inhibiting, controlling or regressing angiogenesis in a human comprising administering over a period of time, a non-toxic dosage amount of a pharmaceutical composition to a human a therapeutically effective non-toxic dosage amount of a composition comprising a non-steroidal anti-inflammatory agent (NSAID), and an effective non-toxic dosage amount of hyaluronic acid and/or salts thereof and/or homologues, analogues, derivatives, complexes, esters, fragments, and/or sub-units of hyaluronic acid which together are effective when the composition is administered to inhibit, control or regress angiogenesis in a human, the said composition further comprising as required pharmaceutical excipients suitable for said form of administration.

25. The method of Claim 24 wherein the pharmaceutical composition is for sub-retinal neovascularisation.

26. The method of Claim 24 wherein the pharmaceutical composition is for the treatment of arthritis or pannus.

27. The method of Claim 24 wherein the pharmaceutical composition is for the treatment of tumours.

28. The method of Claim 24 wherein the pharmaceutical composition is for an adjunct to cancer treatment.

29. The method of Claim 24, 25, 26, 27 or 28 wherein the form of hyaluronic acid is sodium hyaluronate and the molecular weight is less than about 750,000 daltons.

30. The method of Claim 24, 25, 26, 27 or 28 wherein the dosage amount is for systemic administration.

31. The method of Claim 30 wherein the amount of the sodium hyaluronate exceeds about 200 mg/70 kg person and the NSAID is between about 15 mg to about 100 mg of NSAID..

32. The method of Claim 25, 26, 27, 28, 29 or 30 wherein the non-steroidal anti-inflammatory drug (NSAID) is selected from diclofenac sodium in an amount between about 15 mg to about 100 mg/70 kg person and the amount of the form of hyaluronic acid present is 50 mg to 1050 mg.

33. The method of Claim 24, 25, 26, 27, 28, 29, 30, 31 or 32 wherein the NSAID is selected from diclofenac, diclofenac sodium, indomethacin, naproxen, (+/-) tromethamine salt of ketorolac, IBUPROFEN, PIROXICAM, Propionic Acid derivatives, acetylsalicylic acid and Flunixin.

34. The method of Claim 24, 25, 26, 27, 28 or 29 wherein the amount of NSAID in the dosage amount is about 400 mg and the amount of the form of hyaluronic acid is in excess of about 2000 mg.

35. A container containing a composition for administration to a human for the inhibition, control and regression of angiogenesis in humans, the composition comprising at least one dosage amount, each such dosage amount comprising an NSAID and (b) hyaluronic acid and/or sodium hyaluronate, each dosage amount comprising an effective non-toxic dosage amount of each of the NSAID and hyaluronic acid and/or sodium hyaluronate to inhibit, control and regress angiogenesis in a human and wherein the effective amount of the hyaluronic acid and/or sodium hyaluronate comprising about 50 mg of the form of hyaluronic acid for each is about 15 mg of NSAID in each dosage amount to be administered.

36. The container of Claim 35 wherein the hyaluronic acid and/or sodium hyaluronate has a molecular weight less than about 750,000 daltons.

37. The container of Claim 35 or 36 wherein the pharmaceutical composition is for systemic administration.

38. The use of an effective dosage amount of a pharmaceutical composition comprising an effective amount of an NSAID and an effective non-toxic amount of hyaluronic acid and/or salts having a molecular weight less than about 750,000 daltons to inhibit, control and/or regress angiogenesis in a mammal.

39. The use of Claim 38 wherein the pharmaceutical composition is for systemic administration.

40. The use of Claim 39 wherein the amount of the form of hyaluronic acid exceeds about 200 mg/70 kg person and the NSAID is between about 15 mg to about 100 mg of NSAID.

41. The use of hyaluronic acid and/or salts thereof and/or homologues, analogues, derivatives, complexes, esters, fragments, and subunits of hyaluronic acid in the manufacture of a pharmaceutical composition for inhibiting, controlling and/or regressing angiogenesis in a therapy wherein dosage amounts taken from the composition each comprise a therapeutically effective amount of hyaluronic acid and/or salts thereof and/or homologues, analogues, derivatives, complexes, esters, fragments, and sub-units of hyaluronic acid, the pharmaceutical composition being characterized in that for each dosage amount taken from the pharmaceutical composition, the amount of the form of hyaluronic acid inhibits, controls and/or regresses angiogenesis.

42. The use of Claim 41 wherein the pharmaceutical composition is for sub-retinal neovascularisation.

43. The use of Claim 41 wherein the pharmaceutical composition is for the treatment of arthritis or pannus.
44. The use of Claim 41 wherein the pharmaceutical composition is for the treatment of tumours.
45. The use of Claim 41 wherein the pharmaceutical composition is for an adjunct to cancer treatment.
46. The use of Claim 41, 42, 43, 44 or 45 wherein the form of hyaluronic acid is hyaluronic acid and/or salts thereof having a molecular weight less than about 750,000 daltons.
47. The use of Claim 41, 42, 43, 44, 45 or 46 wherein the pharmaceutical composition is for systemic administration.
48. A method of inhibiting, controlling and/or regressing angiogenesis in a human comprising administering a non-toxic dosage amount of a composition comprising a therapeutically effective non-toxic dosage amount of hyaluronic acid and/or salts thereof and/or homologues, analogues, derivatives, complexes, esters, fragments, and/or sub-units of hyaluronic acid which is effective when the composition is administered to inhibit, control and/or regress angiogenesis in a human.
49. The method of Claim 48 wherein the form of hyaluronic acid is sodium hyaluronate having a molecular weight less than about 750,000 daltons.
50. The method of Claim 47 wherein the treatment is administered daily for a number of weeks.
51. The method of treatment of Claim 24 or 48 wherein the treatment comprises administering effective dosage amounts of the composition, a number of times daily for a

period of weeks to inhibit, control and/or regress angiogenesis.

52. The method of Claim 24, 25, 48 or 49 in combination with any method of treating cancer.

53. The use of an effective dosage amount of a composition comprising an effective dosage amount of hyaluronic acid and/or salts thereof having a molecular weight less than 750,000 daltons, sufficient upon administration to inhibit, control or regress angiogenesis in a human.

54. The use of Claim 53 wherein the pharmaceutical composition is for systemic administration.

55. A pharmaceutical composition to inhibit, control or regress angiogenesis in humans, the pharmaceutical composition comprising:

(1) a non-steroidal anti-inflammatory agent (NSAID), and

(2) hyaluronic acid and/or salts thereof and/or homologues, analogues, derivatives, complexes, esters, fragments, and sub-units of hyaluronic acid, in a form suitable for administration to humans; characterized in that said composition comprises at least one effective dosage amount of components (1) and (2) to inhibit, control or regress angiogenesis.

56. The pharmaceutical composition of Claim 55 wherein at least one effective dosage amount comprises a plurality of dosage amounts.

57. The pharmaceutical composition of Claim 55 wherein at least one effective dosage amount comprises one dosage amount.

58. The pharmaceutical composition of Claim 55, 56, or 57 wherein the hyaluronic acid and/or salts thereof

and/or homologues, analogues, derivatives, complexes, esters, fragments, and/or sub-units of hyaluronic acid is hyaluronic acid and/or a salt thereof having a molecular weight less than about 750,000 daltons.

59. The pharmaceutical composition of Claim 55, 56, 57 or 58 wherein the form of hyaluronic acid is sodium hyaluronate having a molecular weight less than 750,000 daltons and the sodium hyaluronate in each dosage amount is present in an amount of about 2200 mg and the diclofenac is present in an amount of about 420 mg.

60. The use of Claim 38 or 53 for the treatment of sub-retinal neovascularisation.

61. The use of Claim 38 or 53 for the treatment of arthritis or pannus.

62. The use of Claim 38 or 53 for the treatment of tumours.

63. The use of Claim 38 or 53 as an adjunct to cancer treatment.

64. The method of Claim 24, 25, 26, 27, 28 or 29 wherein the form of hyaluronic acid is sodium hyaluronate having a molecular weight less than 750,000 daltons and the sodium hyaluronate in each dosage amount is present in an amount of about 2200 mg and the diclofenac is present in an amount of about 420 mg.




FIG.1A.




FIG.1B.




FIG. 2.

3 / 4

Anti-Angiogenesis: Carmine




FIG. 3.

4 / 4

Anti-Angiogenesis:
Dry Weight Granuloma




FIG.4.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 94/00207

A. CLASSIFICATION OF SUBJECT MATTER
IPC 5 A61K31/725

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 433 817 (NISSHO CORPORATION) 8 December 1990 see the whole document ---	1-64
X	WO,A,91 04058 (NORPHARMCO INC.) 4 April 1991 see claims 2,3 ---	1-64
X,P	WO,A,93 16733 (NORPHARMCO INC.) 2 September 1993 see claims 1,7,10 -----	1-64

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

22 July 1994

Date of mailing of the international search report

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Leherte, C

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 94/00207

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0433817	26-06-91	DE-D- 69003229	14-10-93
		DE-T- 69003229	14-04-94
		JP-A- 3236330	22-10-91
		US-A- 5095037	10-03-92

WO-A-9104058	04-04-91	AU-B- 5227493	03-03-94
		AU-A- 6433090	18-04-91
		CN-A- 1051503	22-05-91
		EP-A- 0445255	11-09-91
		JP-T- 4504579	13-08-92

WO-A-9316733	02-09-93	CA-A- 2061566	21-08-93
		AU-B- 3488993	13-09-93

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA94/00207

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
REMARK: Although claims 24-34, 38-54, 60-64 are directed to a method of treatment of (diagnostic method practised on) the human/animal body the search has been carried out and based on the alleged effects of the compound/
COMPOSITION.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.